



NR 2:1 BEEF CATTLE PREMIX MONENSIN WITH SALT

Medicated with 3,300 mg/kg monensin sodium for improved feed efficiency in beef cattle (steers and heifers) fed in confinement for slaughter. A list of the ingredients used in this feed may be obtained from the manufacturer. This feed contains added selenium at 30.0 mg/kg.

GUARANTEED ANALYSIS:

Calcium, Act.....	16.00 %
Phosphorus, Act.....	7.50 %
Magnesium, Act.....	2.00 %
Sodium, Act.....	6.00 %
Salt, Act	15.24 %
Sulfur, Act.....	0.15 %
Copper, Act.....	2,500 mg/kg
Zinc, Act.....	7,000 mg/kg
Manganese, Act.....	5,000 mg/kg
Iron, Act.....	3,500 mg/kg
Iodine, Act.....	100 mg/kg
Cobalt, Act.....	35 mg/kg
Vitamin A, Min.....	500,000 iu/kg
Vitamin D, Min.....	60,000 iu/kg
Vitamin E, Min.....	2,000 iu/kg
Flourine, Max	900 mg/kg

FEEDING DIRECTIONS:

Thoroughly mix this premix into the total daily beef cattle diet or in complete feed (grain portion of the ration) before use. Do not feed undiluted. For example, mix 10 kg of this premix along with 2 kg of Dyna K, 16 kg of Calcium and 972 kg of barley or mixed grain to make one tonne of beef cattle ration. Contact a New Rosedale Feedmill representative for more information. Feed the resulting ration to finishing beef cattle according to the table below (based on a Dry Matter Intake (DMI) of 2.5% body weight):

BODY WEIGHT (BW)	D.M.I. AS % OF BW	MIXED FEED INTAKE KG/DAY	DAILY INTAKE OF MONENSIN ACTIVITY (MG/HEAD/DAY)
250	2.50	6.25	206
300	2.50	7.50	248
350	2.50	8.75	289
400	2.50	10.00	330

WARNING:

1. Do not supplement monensin from other sources (e.g., other feedstuffs containing monensin or slow release devices containing monensin).
2. No withdrawal period or milk withholding time is required when treated up to 48 mg monensin per kg (0.0048%) of complete diet.
3. When mixing and handling monensin, avoid inhalation, oral exposure and direct contact with skin or eyes. Use protective clothing, impervious gloves and dust mask. Operators should wash thoroughly with soap and water after handling.
4. Keep out of reach of children.

CAUTION:

1. Directions for use must be carefully followed.
2. Do not use in association with another feed containing supplemental selenium.
3. Feed only to beef cattle (steers and heifers) intended for slaughter and maintained under confinement.
4. Do not exceed recommended levels as reduced average daily gains may result.
5. Do not allow dogs, horses, other equines or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.
6. May be used in feeds containing the pellet-binding agents Bentonite (2%), Attapulgit (2%), Kaolin (2.5%), Lignin Sulfonate (4%), Carboxymethyl cellulose (0.1%) or Agri-Colloid.

Individual results from the use of this product may vary due to management, environmental, genetic, health and sanitation differences; therefore, New Rosedale Feed Mill does not guarantee individual results.

Manufactured by:
NEW ROSEDALE FEED MILL
PORTAGE LA PRAIRIE, MANITOBA

Net weight: bulk or 25 kg when bagged formula

Code 4186_